https://invertebratefungi.org/: an expert-curated web-based platform for the identification and classification of invertebrate-associated fungi and fungus-like organisms

De-Ping Wei^{1,2,3}, Eleni Gentekaki^{2,4}, Kevin D. Hyde^{1,2,5,8,*}, Yuan-Pin Xiao^{2,4}, Thatsanee Luangharn², Qing Tian^{2,4}, Yuan-Bing Wang⁶ and Saisamorn Lumyong^{7,8,9}

¹CAS Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Panlong District, Kunming, Yunnan, P.R. 650201, China

²Center of Excellence in Fungal Research (CEFR), Mae Fah Luang University, Muang District, Chiang Rai 57100, Thailand
³Department of Entomology and Plant Pathology, Faculty of Agriculture, Chiang Mai University, Mueang Chiang Mai, Chiang Mai 50200, Thailand

⁴School of Science, Mae Fah Luang University, Muang District, Chiang Rai 57100, Thailand

⁵Innovative Institute of Plant Health, Zhongkai University of Agriculture and Engineering, Haizhu District, Guangzhou 510225, P.R. China
⁶The International Joint Research Center for Sustainable Utilization of Cordyceps Bioresources in China and Southeast Asia, Yunnan University, Wuhua District, Yunnan, P.R. 650091, China

⁷Department of Biology, Faculty of Science, Chiang Mai University, Mueang Chiang Mai, Chiang Mai 50200, Thailand

⁸Research Center of Microbial Diversity and Sustainable Utilization, Faculty of Science, Chiang Mai University, Mueang Chiang Mai, Chiang Mai 50200, Thailand

⁹Academy of Science, The Royal Society of Thailand, Dusit District, Bangkok 10300, Thailand

*Corresponding author: Tel: +66 53916961; Email: kdhyde3@gmail.com

Citation details: Wei, D., Gentekaki, E., Hyde, K.D. *et al.* https://invertebratefungi.org/: an expert-curated web-based platform for the identification and classification of invertebrate-associated fungi and fungus-like organisms. *Database* (2022) Vol. 2022: article ID baac021; DOI: https://doi.org/10.1093/database/baac021

Abstract

Fungi are the major decomposers in terrestrial and aquatic ecosystems, playing essential roles in biogeochemical cycles and food webs. The Fungi kingdom encompasses a diverse array of taxa that often form intimate relationships with other organisms, including plants, insects, algae, cyanobacteria and even other fungi. Fungal parasites of insects are known as entomopathogenic fungi and are the causative agents of serious disease and/or mortality of their hosts. Entomopathogens produce distinct metabolic compounds with roles in pathogenicity, virulence and host-parasite interactions. Thus, the potential of discovering new bioactive compounds useful in biocontrol and pharmaceutical industries is high. Given the significance of entomopathogenic fungi, the rapid research advances and the increased interest, it has become necessary to organize all available and incoming data. The website https://invertebratefungi.org/ has been developed to serve this purpose by gathering and updating entomopathogenic genera/species information. Notes of entomopathogenic genera will be provided with emphasis on their taxonomic status. Information on other invertebrates, such as rotifers, will also be included. Descriptions, photographic plates, information on distribution and host (where applicable) along with molecular data and other interesting details will also be provided. The website is easily and freely accessible to users. Instructions concerning the platform architecture and functionality of the website are introduced herein. The platform is currently being expanded and will be continuously updated as part of the effort to enrich knowledge on this group of fungi.

Database URL: https://invertebratefungi.org/

Introduction

Invertebrate organisms account for a major part of animal diversity. Over 97% of all described animal species are invertebrates being omnipresent in terrestrial and aquatic ecosystems (1). Fungi have the ability to colonize a wide range of invertebrate hosts, ranging from soil-inhabiting microinvertebrates to macroinvertebrates in Insecta. Fungi can live as entomopathogens, either endoparasites or ectoparasites, in order to exploit their invertebrate hosts and obtain nutrients (7–11). Invertebrates are known to harbor a vast number of fungi and fungus-like organisms (2). The most available

literature on invertebrate fungi focuses on entomopathogens. The taxonomic framework of entomopathogenic fungi has undergone several revisions (2-6). Phylogenetic analyses have shown that entomopathogenic fungi are distributed across six major groups: Ascomycota, Basidiomycota, Chytridiomycota, Microsporidia, zygomycetes and fungus-like Oomycota (2, 5, 6).

The phylum Ascomycota contains high diversity of entomopathogenic fungi, which mainly parasite arthropod hosts in forests and soil-dwelling microinvertebrates such as nematodes and rotifers (9, 12-16). All members of the order

Received 5 January 2022; Revised 17 February 2022; Accepted 21 March 2022

© The Author(s) 2022. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Contents presented under eight main headings



Figure 1. A schematic of the database structure.

Laboulbeniales (Ascomycota) are obligate ectoparasites of the integument of insects, but the effect on their hosts is not lethal (17, 18). Entomopathogenic Basidiomycota comprises the termite egg-infecting genus Fibularhizoctonia and the scale insect-parasitic genera Uredinella and Septobasidium (2). Chytridiomycota includes the entomopathogenic genera Myiophagus, Coelomycidium, Myrmicinosporidium and Coelomomyces (2). These infect mostly Diptera hosts and occasionally scale insects (2). Microsporidia is a group of intracellular pathogenic fungi, consisting of 143 genera, 69 of which are known to infect a broad array of insect hosts spanning 12 orders (2). Zygomycetes along with Trichomycetes were initially accommodated in the phylum Zygomycota (19). However, recent phylogenetic analyses have revealed the polyphyletic nature of Zygomycota and, as a consequence, this phylum has now been abandoned and replaced with multiple new phyla (20-26). Entomophthoromycota and

Mucoromycota are two of the newly erected Zygomycetes phyla containing entomopathogenic fungi. In Entomophthoromycota, entomopathogens are distributed across 19 genera forming biotrophic associations across 10 orders of insects (2). Mucoromycota contains a large number of Trichomycetes fungi, which are obligate symbionts of the gut of various arthropod larvae in aquatic environments (2). Several genera in Oomycota, namely *Aphanomyces, Couchia, Crypticola, Lagenidium, Leptolegnia* and *Pythium*, have been recorded as pathogens of mosquito, midge, lobster and shrimp (2, 27–29).

Invertebrate-associated fungi are highly diverse, with entomopathogens comprising a major constituent of this group. Entomopathogenic fungi play important roles in keeping populations of insects under control in nature and thus have vast potential as biocontrol agents. They also constitute a rich resource of naturally produced bioactive compounds. Many



Figure 3. The outline of invertebrate-associated fungi and fungi-like organisms.

biological insecticides have been developed from this group of fungi and used in protecting crops from pests (30–32). A large variety of secondary metabolites produced by entomopathogenic fungi have been validated as nutraceuticals and traditional remedies (33). However, only a fraction of entomopathogenic species has been successfully exploited for their biocontrol and pharmaceutical value potential. Informal identification or improper taxonomic placement of these fungi is a frequent issue due to the dynamic nature of fungal classification system. This may lead to increasing taxonomic confusion among scientists and mycologists. Therefore, it is essential to provide their corrected fungal names accompanied by information about the biology, host range, distribution and specific functions (34). The accurate identification and precise scientific documentation of these fungi not only benefits communication with mycologists and interested amateurs

10011												
Invertebrate Fungi Heirarch	iy								Searc	h		
Ascosphaerales												
Capnodiales									Recen	nt Genus		
Hypocreales - Order							Stigmatomyces Metarbizionsis					
Read more about Hypocreales orders »									Metapo	ochonia		
Clavicipitaceae									Recen	nt Species		
Cordycipitaceae									Ophioc	ordyceps kars	tii	
Ophiocordycipitaceae - Famil	у								Ophioc	tordyceps forn tomyces vikhr	nosana evii	
Read more about Ophiocordycipitacea	e family »											
Drechmeria												
Ophiocordyceps		10										
Read more about Ophiocordyceps g	enus » 🔶	Genus										
Ophiocordyceps cossidarum												
Ophiocordyceps cylindrospora												
Ophiocordyceps formosana	← Sp	ecies										
Ophiocordyceps issidarum												
Ophiocordyceps karstii												
4. The 'Archives' section ext	FUN	ngal taxa G I	in hierarchi	cal order.	Home	Outline	Archives	Curators	History	References	News	Co
4. The 'Archives' section ext	FUN	ngal taxa G I	in hierarchi	cal order.	Home	Outline	Archives	Curators	History	References	News	Co
4. The 'Archives' section exit NVERTEBRATE SSN ***********************************	ribiting fur	ngal taxa	in hierarchi 1) iocordycipitacea cies (Species Fi	cal order.	Home 125 species	Outline with seque	Archives	Curators	History Rece Stigm Metar Metar	References ent Genus atomyces hiziopsis pochonia	News	Co
4. The 'Archives' section ext NVERTEBRATE SSN ***********************************	FUN vcol. Soc. 16(tidae, Hypocr ; 242 morpho e. Sexual mo	ngal taxa G 1): 73 (193 eales, Oph logical spe rph: Stron	in hierarchi 1) locordycipitacee cies (Species Fi aata or subiculu	cal order. ae ungorum 2020), m darkly pigmen	Home 125 species ited or rarely	Outline with seque	Archives ence data.	Curators	History Rece Stigm Metar Metap	References ent Genus atomyces hiziopsis pochonia ent Species	News	Со
4. The 'Archives' section exh NVERTEBRATE SSN ***********************************	ribiting fur FUN ycol. Soc. 16(tidae, Hypocr ; 242 morpho e, Sexual mo e, Sexual mo d, rarely fusoi	ngal taxa G 1): 73 (193 eales, Oph logical spe orph: Strom pices or lat d to ellipso	in hierarchi 1) iocordycipitacea cies (Species Fi nata or subiculu eral pads. Asco id, usually with	cal order. ae ungorum 2020), <i>m</i> darkly pigmen mata superficial thickened ascus	Home 125 species hted or rarely to complete apex. Ascos	Outline with seque brightly o by immerse	Archives ence data. Joured, toug d, ordinal ou iculate, hyai	Curators gh, fibrous, r oblique in line usually	History Rece Stigm Metar Metag Rece Ophic	References ent Genus atomyces rhiziopsis pochonia ent Species pocordyceps kar	News	Co
4. The 'Archives' section exh NVERTEBRATE SSN typocreales » Ophiocordycipitaceae Ophiocordyceps Dphiocordyceps Petch, Trans. Br. my Sordariomycetes, Hypocreomyce Index Fungorum number: IF 3598 Parasitic on arthropods, worldwid dilant to wiry, rarely fleshy, often with a tranagement. Asci cylindrical, 8-spore- ylindrical, multi-septate, disarticulatin Type species: Ophiocordyceps b Notes: Ophiocordyceps is the typ in morphology and phylogenetic an Sorosporella, a synonym of Synglioc reased with respect to the second ol ames. Most species of the genus aru resselie etomate europricel to co-	vcol. Soc. 16(tidae, Hypocr ; 242 morpho e, Sexual mo aperthecial a d, rarely fusoi g into part-sp lattae (Petch) alyses. Base cladium. Soro dest name H e considered	1): 73 (193 eales, Oph logical spe orph: Stron pices or lat d to ellipso phiocordycd d on phyl isporella w irsutella (E as entomo	in hierarchi 1) iocordycipitacea cies (Species F nata or subiculu eral pads. Asco ueral pads. Asco ipitaceae (Sung ogenetic analys as suppressed vans & Samson pathogenic and thoria and asco	cal order. ae ungorum 2020), m darkly pigmen mata superficial inickened ascus (Maharachchikur y et al. 2007). Qu ses, the oldest (Quandt et al. n 1982). <i>Ophioc</i> have darkly pig learther have darkly pig learther	Home 125 species ted or rarely to complete apex. Ascox mbura et al. (in ame in th 2014), while cordyceps sh mented, touy usting into	Outline with seque brightly co brightly co brightly co brightly co brightly co co co brightly co brightly co co co brightly co brightly co co co brightly co co co co co co co co co co co co co c	Archives Archives ance data. Joured, toug d, ordinal ou iculate, hyai anded this ge adjum was otected ove pliant to wi or pon-di	Curators gh, fibrous, r oblique in line usually enus based e clade is previously r the other ry, stipitate adiculatios	History Rece Stigm Metar Metap Rece Ophic Ophic Stigm	References ent Genus atomyces rhiziopsis pochonia ent Species pocordyceps kai pocordyceps for atomyces vikh	News rstii rmosana irevii	Co
4. The 'Archives' section exh NVERTEBRATE SSN typocreales » Ophiocordycipitaceae Ophiocordyceps Dphiocordyceps Petch, Trans. Br. m; Sordariomycetes, Hypocreomyce Index Fungorum number. IF 3598 Parasitic on arthropods, worldwid iliant to wiry, rarely fleshy, often with a transgement. Asci cylindrical, 8-spore ylindrical, multi-septate, disarticulatin Type species: Ophiocordyceps b Notes: Ophiocordyceps is the typ in morphology and phylogenetic an Sorosporella, a synonym of Synglioc reated with respect to the second of names. Most species of the genus aru r sessile stromata, superficial to co Sung et al. 2007). The asexual morpho 1007). (Adopted from Hyde et al. 2020 Host range: Lepidoptera, Hemip	recol. Soc. 16(tidae, Hypocr ; 242 morpho aperithecial a d, rarely fusoi g into part-sp lattae (Petch) lattae (Petch) cladium. Sorc dest name H e considered e considered mpletely imm ns of this gen)) tera, Hymeno	ngal taxa () 1): 73 (193 eales, Oph logical spe orph: Stron pices or non Petch phiocordycd d on phyl isporella w irsutella (E as entomo ersed peri us are mer optera.	in hierarchi 1) iocordycipitacea cies (Species Fi- nata or subiculu eral pads. Asco d, usually with -disarticulating ipitaceae (Sung ogenetic analy as suppressed vans & Samson pathogenic and thecia and asco nbers of Hirsufe	cal order. ae ungorum 2020), m darkily pigmen mata superficial thickened ascus (Maharachchikur y et al. 2007). Qu yes, the oldest (Quandt et al. n 1982). Ophioc have darkly pig pospores disartici lila, Hymenostibi	Home 125 species hed or rarely to complete apex. Ascox mbura et al. (in ame in th 2014), while cordyceps sh mented, touy ulating into p le, Paraisaria	Outline brighty or by immerse pores faso 2016). 2014) eme e Ophioco a Synglioc ould be pr gh, fibrous part-spores a and Syng	Archives Arc	Curators gh, fibrous, r oblique in line usually enus based he clade is previously r the other ry, stipitate articulating Sung et al.	History Rece Stigm Metar Metap Rece Ophic Ophic Stigm	References atomyces rhiziopsis pochonia atomyceps kar bcordyceps kar bcordyceps kar bcordyceps vikh	News	Co
4. The 'Archives' section exit NVERTEBRATE SNVERTEBRATE SNVERTEBRATE SN typocreales » Ophiocordycipitaceae Ophiocordyceps Dohiocordyceps Dohiocordyceps Petch, Trans. Br. my Sordariomycetes, Hypocreomyce Index Fungorum number. IF 3598 Parasitic on arthropods, worldwid diiant to wiry, rarely fleshy, often with a irrangement. Asci cylindrical, 8-spore ylindrical, multi-septate, disarticulatin Type species: Ophiocordyceps b Notes: Ophiocordyceps is the typ in morphology and phylogenetic an Sorosporella, a synonym of Synglice reated with respect to the second of ames. Most species of the genus and r sessile stromata, superficial to co Sung et al. 2007). The asexual morpli 007). (Adopted from Hyde et al. 2020 Host range: Lepidoptera, Hemip Dther species: Dahogordyceps cossidarum V.P. Xiao	real provided and the second s	1): 73 (193 eales, Oph logical spe orph: Stron pices or lat d to ellipso prices or non Petch phiccordyc d on phyl sysperilla w irsutella (E as entomo ersed peri us are mer optera.	in hierarchi 1) locordycipitacea cies (Species Fi nata or subiculu eral pads. Asco id, usually with -disarticulating ipitaceae (Sung ogenetic analys as suppressed vans & Samson pathogenic and thecia and asco nbers of <i>Hirsute</i> tyde	ae ungorum 2020), m darkly pigmen wata superficial thickened ascus (Maharachchikur et al. 2007). Qu ses, the oldest (Quandt et al. n 1982). Ophioc have darkly pig ospores disarticu	Home 125 species ted or rarely to complete apex. Ascoc mbura et al. (name in the Jordy ceps sh mented, tougulating into le, Paraisaria	Outline with seque brightly c ly immerses 2016). 2014) eme e Ophioco a Synglioc ould be pr ph, fibrous part-spores a and Syng	Archives ence data. bloured, toug d, ordinal or iculate, hyai nded this ge rdycipitacea adium was otected ove pliant to wi or non-dis liocladium (Curators gh, fibrous, r oblique in line usually enus based ec clade is previously r the other rty, stipitate articulating Sung et al.	History Rece Stigm Metar Metar Ophic Ophic Stigm	References atomyces hiziopsis bochonia atot Species bocordyceps kar bocordyceps for atomyces vikh	News rstii mosana nrevii	Co
4. The 'Archives' section exit NVERTEBRATE SNVERTEBRATE SNVERTEBRATE SNVERTEBRATE SNVERTEBRATE SNVERTEBRATE SNVERTEBRATE SNT Apported to the sector of the sector	rool. Soc. 16(tidae, Hypocr ; 242 morpho e. Sexual mo aperithecial a d, rarely fusoi g into part-sp earch fusoi g into part-sp lattae (Petch) e genus of O alyses. Base cladium. Sorc dest name H e considered mpletely imm is of this gen)) tera, Hymeno ; T.C. Wen, Z & Shimizu) V Xiao & K.D. H	1): 73 (193 eales, Oph logical spe orph: Stroon Petch phiccordyc d on phyl isporella w irsutella (E as entomo ersed peri us are mer optera. ha & K.D. Hyd Vang et al. yde [as 'ka	in hierarchi 1) iocordycipitacea cies (Species Fi- hata or subiculu real pads. Asco- id, usually with -disarticulating ipitaceae (Sung ogenetic analys as suppressed yans & Samso- nata or <i>Hirsute</i> e in Wang et al. still), in Li et al.	ee ungorum 2020), m darkly pigmen mata superficial thickened ascus (Maharachchikur u et al. 2007). Qu ses, the oldest (Quandt et al. (Quandt et al. (Quandt et al. n have darkly pig ospores disarticu bila, Hymenostibil	Home 125 species ted or rarely to complete apex. Ascos mbura et al. (andt et al. (name in th 2014), while ordyceps sh mented, tougulating into le, Paraisaria 3: Click	Outline with seque brightly of y immerse pores faso 2016). 2014) eme e Ophioco a Synglioc ould be pr jh, fibrous part-spores a and Syng	Archives Arc	Curators gh, fibrous, r oblique in line usually enus based e clade is previously r the other ry, stipitate Sung et al.	History Rece Stigm Metar Metar Rece Ophic Stigm	References atomyces hiziopsis pochonia ent Species pocordyceps kar pocordyceps for atomyces vikh	News rstii rmosana irevii	Con
4. The 'Archives' section exh NVERTEBRATE SNVERTEBRATE SNVERTEBRATE SNVERTEBRATE SNVERTEBRATE SNATTANA Apported to the sector of the se	FUN FUN (col. Soc. 16() (col. Soc. 16() (co	1): 73 (193 eales, Oph logical spe orph: Stron pices or lat d to ellipso cores or non Petch phiccordyc d on phyl isporella w irsutella (E as entomo iersed peri us are mer optera. ha & K.D. H & K.D. Hya Vang et al. yde [as 'ka & K.D. Hy	in hierarchi 1) iocordycipitacea cies (Species Fi nata or subiculu eral pads. Asco- id, usually with -disarticulating ipitaceae (Sung ogenetic analys as suppressed vans & Samson pathogenic and thecia and asco- nbers of <i>Hirsute</i> tyde e in Wang et al	ee ungorum 2020), m darkly pigmen mata superficial thickened ascus (Maharachchikur of at. 2007). Qu ses, the oldest (Quandt et al. n 1982). Ophioc have darkly pig ospores disarticu alla, Hymenostibi	Home 125 species ted or rarely to complete apex. Ascoss mbura et al. (name in th 2014), while ord/cops sh mented, touy ulating into le, Paraisaria 3: Click	Outline with seque brightly of ly immerse pores fasc 2016). 2014) eme e Ophioco a Synglico ould be pr jh, fibrous part-spores a and Syng	Archives ance data. Joured, toug d, ordinal ou iculate, hyai adium was otected ove pliant to wi or non-dis <i>liocladium</i> (Curators gh, fibrous, r oblique in line usually enus based e clade is previously r the other ry, stipitate articulating Sung et al.	History Rece Stigm Metar Metar Ophic Ophic Stigm	References atomyces hiziopsis bochonia atotyceps kar bocordyceps kar bocordyceps vikh	News rstii mosana nrevii	Co

but also paves the way for downstream utilization, including pharmaceutical industry and biological control applications.

To accumulate and organize available data in a single repository and to effectively communicate the knowledge of fungi to the public, many databases have been designed for different fungal groups, such as Basidiomycota (35), Botryosphaeriales (36), Coelomycetes (https://www.coelo

mycetes.org/), Dothideomycetes (37), fresh water fungi (38), phytopathogens (39) and Sordariomycetes (40). In addition, notes on changes and additions to phyla, classes, orders, families and genera of fungi are provided on the website Outlineoffungi.org (https://www.outlineoffungi.org/). However, a database dedicated to fungi and fungi-like organisms associated with invertebrates has not yet been developed. INVERTEBRATE FUNGI ISSN ****-**** Hypocreales » Ophiocordycipitaceae » Ophiocordyceps

Ophiocordyceps formosana

Ophiocordyceps formosana (Kobayasi & Shimizu) Wang et al. in Wang et al., Evidence-Based Complementary and Alternative Medicine (no. 189891): 4 (2015)

Index Fungorum Number – IF 811035; Facesoffungi Number: FoF 01796 +FOF number

Parasitic in larva of Coleoptera (Superfamily Tenebrionoidea), forming yellow to orange ascostromata. Sexual morph: Ascomycetous. Stromata 14 mm long, 2-5 mm wide, growing from the head and the tail of Coleoptera larva, simple or branched, yellow to orange, stipitate. Stipe 1.9–3.7 cm long, 2–4 cm wide, yellow, cylindrical, surface rough. Fertile head 30 mm long, 2–2.5 mm wide, orange, mostly elliptic barely branched, head-like, with orange, pseudoparenchymatous, epidermal tissues, surface mastoid, differentiated from stipe. Ascomata 453–546 × 265–298 µm ($\hat{x} = 479 \times 270$, n = 30), completely immersed, orange, flask-shaped to oval, with the ostioles opening on the surface of the head. Peridium 26–38 µm wide ($\hat{x} = 30$, n = 60), comprising three layers. Asci 366–498× 8–11 µm ($\hat{x} = 437 \times 10$, n = 60), 8-spored, hyaline, cylindrical, with apical cap, breaking into secondly ascospores. Secondary ascospores 2–6 × 1–3 µm ($\hat{x} = 4 \times 2$, n = 60), hyaline, cylindrical. Asexual morph: Undetermined. (Adopted from Li et al. 2016) Recent Genus Stigmatomyces

Metarhiziopsis Metapochonia

Recent Species

Ophiocordyceps karstii Ophiocordyceps formosana Stigmatomyces vikhrevii

- Species description

Known hosts and distribution: on dead larva of Tenebrionoidea (China) (Li et al. 2016) Moelecular data: MFLU 15-3888 (SSU = KU854951, RPB1 = KU854947, TEF = KU854949). Phylogenetic tree: see entry (Ophiocordyceps karstii) (Li et al. 2016) Molecular data



Figure 6. Information of the entomopathogenic species Ophiocordyceps formosana.

The website invertebratefungi.org is therefore established as part of the project titled 'The future of specialist fungi in a changing climate: baseline data for generalist and specialist fungi associated with ants, *Rhododendron* species and *Dracaena* species'. The aims of this website are to: (i) collect and curate morphological and molecular data on invertebrateassociated fungi from formally published articles and (ii) provide a continuously updated checklist of entomopathogenic genera/species.

The need for a database of invertebrate fungi

In the past decade, the application of molecular tools has led to a dramatic revolution in the taxonomy of fungal entomopathogens. Molecular analyses have revealed several cases where phenotypic features do not necessarily reflect phylogenetic results (12, 14). A distinct example comes from clavicipitaceous fungi. Several subgenera under Cordyceps sensu lato (Clavicipitaceae) were initially recognized on the basis of the arrangement of perithecia, the disarticulation of ascospores and host affiliation (41–44).

Later works revealed that the aforementioned characteristics presented cases of convergent evolution in Clavipitaceae sensu lato (12). Consequently, Clavicipitaceae sensu stricto, Cordycipitaceae and Ophiocordycipitaceae were erected to accommodate species of Cordyceps s. l. and other clavicipitaceous fungi. Combination of morphological and phylogenetic analyses has been crucial for resolving numerous taxonomic issues and for providing a reliable, more robust taxonomic framework of entomopathogenic fungi (12, 45). However, many species that were collected and described before the molecular era lack molecular data, which poses a significant challenge to their natural classification. Fresh collections of these species and clear morphological illustrations would aid tremendously in clarifying their naturally taxonomic placements, and these are currently being undertaken. Although many newly recorded and novel fungal species on invertebrates have been discovered, described and published across a plethora of journals, the published information is scattered and often not easily accessible. Moreover, the classification of entomopathogenic fungi is constantly updated, and these changes are hard to follow. Comprehensively compiling the

ISSN ****-***							
Туј	be the order name	of ins	sect hos	sts			
Genus: Cordyceps					Lepio	doptera	
ented, fleshy, stromata growing from the heads of adult Lepidoptera, usually s	simple, rarely branched More				Dece	nt Nours	
Genus: Paraisaria					Rece	nt Conuc	
leoptera, Dictyoptera, Diptera, Hemiptera, Hymenoptera, Lepidoptera and Ort	hoptera (Wei et al. 2021) Mo	e			Rece	int Genus	
Genus: Ophiocordyceps							
al. 2007). (Adopted from Hyde et al. 2020) Host range: Lepidoptera, Hemipter	ra, Hymenoptera More						
Species: Cordyceps militaris							
Index Fungorum number: IF 237604 Parasitic on pupa of Lepidoptera. Sexua	al morph: Stroma 2–7 More						
Species: Cordyceps pruinosa							
on cocoon of a Limacodidae insect, Iragoides fasciata (Lepidoptera) (China,	Korea, Japan) (Hyde et More	2					
Species: Cordyceps rosea							
IF 114258, Facesoffungi number: FoF07467 Parasitic on Lepidoptera larva o	r pupa. Sexual morph: Str Mo	re					
Species: Paraisaria arcta							
, Facesoffungi number: FoF 07240 Parasitic on larva of Lepidopteran larva. S	exual morph: Stroma 16 Mor	e					
Species: Ophiocordyceps cossidarum							
fungi number: FoF 3278 Parasitic in larvae of insect (Lepidoptera: Cossidae)	, collected from burie More						

Position	Name	Research field	Contact details		
Head Curator	Yuan-Pin Xiao ^a	Entomopathogenic fungi	emmaypx@gmail.com		
Managing Curator	De-Ping Wei ^a	Entomopathogenic fungi	elina19920601@gmail.com		
Curator	Kevin D. Hyde ^a	Fungi	kdhyde3@gmail.com		
Curator	Qing Tian ^a	Eurotiomycetes	tianging124@gmail.com		
Curator	Yuan-Bing Wang ^b	Entomopathogenic fungi	wangyb001@126.com		

^aCenter of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand.

^bThe International Joint Research Center for Sustainable Utilization of Cordyceps Bioresources in China and Southeast Asia, Yunnan University, Kunming 650091, China.

taxonomic data concerning entomopathogenic fungi according to the original publications is in high demand by peers and researchers in related fields. An easily accessible and freely available platform would greatly facilitate the sharing and dissemination of knowledge among mycologists and other scientists. The online database is dedicated to gathering information from various peer-reviewed sources, allowing users to access up-to-date data in a time-saving and effective way. The website https://invertebratefungi.org/ was therefore established to systematically document data on invertebrate fungi in a web-based platform through which the end users can easily obtain access to relevant information.

Invertebrate fungi website

The website contributes to the checklist of fungal genera and species associated with invertebrates. Current notes regarding taxonomic status, classification, host range and species with available morphological descriptions of fungal genera will be provided. For each genus, the species associated with invertebrate hosts will be presented and accompanied by Index Fungorum number, Facesoffungi number, descriptions, photographic plates, sequence data, phylogenetic data if available, known hosts and distributions, as well as other relevant information of importance. Systematic management and documentation of invertebrate-associated fungi will enhance the understanding of diversity and classification of invertebrate fungal pathogens.

Construction, interface and visualization of database

The online website, https://invertebratefungi.org/, allows end users to enter information on fungi associated with invertebrates via a friendly interface. This website comprises eight headings, each with its own distinct functionality (Figure 1). The research scope of this website is presented at the top-left corner of the homepage (Figure 2). The contact details and supporting grant information are located at the bottom of this page (Figure 2). The genera containing entomopathogenic species have been collected from peer-reviewed articles and systematically listed under the heading 'Outline' according to the most recent classification (Figure 3). Genera/species with available data will be automatically arranged in alphabetical order under the heading 'Archives' (Figure 4). Data of each genus and species presented on the website have been mainly extracted from original research articles published by the Center of Excellence in Fungal Research (CEFR) group. Data from

collaborating institutes will also be included after getting permission. Otherwise, the taxonomic status of the genus will be noted and the entomopathogenic species will be listed along with a hyperlink to the original literature. Researchers who do not belong to the CEFR group also have the option to post the entomopathogenic species they discover via the managing curator of this website. The database will be updated continually to keep abreast of the current literature. Curators listed in the table 1 will take charge of collecting, managing, overseeing and reviewing the uploaded data in this webpage.

How to use the website?

Instructions on searching a specific genus and species are illustrated below using the genus *Ophiocordyceps* and the species *O. formosana* as examples. Step 1: typing a specific genus name (e.g. *Ophiocordyceps*) into the search box located at the top-left corner of the 'outline' page allows users to find the genus in the outline (Figure 3). Step 2: a new window with information on *Ophiocordyceps* will pop up after clicking on the genus name (Figure 5). Step 3: the entomopathogenic species belonging to *Ophiocordyceps* will be presented beneath the 'other species' title, and users can find detailed information on a specific species (e.g. *O. formosana*) by clicking on the species name (Figure 5 and 6). The Facesoffungi Number (Figure 6) of a species is provided along with a hyperlink to the 'Faces of fungi' website (https://www.facesoffungi.org/).

Additionally, a search box is provided below the header bar, allowing the users to search the information on a specific genus and species by typing its name (Figure 2). Recently uploaded genera and species can be seen under 'Recent Genus' and 'Recent Species' tabs, respectively, which inform users of the newly added entries (Figure 2). Alternatively, clicking on the order name in the 'archive' page will result in a dropdown box appearing, where families of the selected order are displayed, and users can browse the genus and species composition of the selected family (Figure 4).

Users can get a list of fungal species associated with specific hosts by typing the scientific or popular name of insect hosts in the search box below the eight header bars (see Figure 7).

Funding

This work was supported by the Chinese Research Fund (project no E1644111K1) entitled 'Flexible introduction of high-level expert program, Kunming institute of botany, Chinese academy of sciences'. The Thailand Research Fund (grant no DBG6080013) entitled 'The future of specialist fungi in a changing climate: baseline data for generalist and specialist fungi associated with ants, *Rhododendron* species and *Dracaena* species' is acknowledged for its financial support. K.D.H. thanks Chiang Mai University for the award of Visiting Professor.

Conflict of interest

All authors have no conflict of interest.

References

1. May,R.M. (1988) How many species are there on earth? *Science*, 241, 1441–1449.

- Araújo, J.P.M. and Hughes, D.P. (2016) Diversity of entomopathogenic fungi. Which groups conquered the insect body? *Adv. Genet.*, 94, 1–39.
- Samson,R.A., Evans,H.C. and Latgé,J.-P. (1988) Taxonomy of entomopathogenic fungi. In: *Atlas of Entomopathogenic Fungi*. Springer, Berlin, Heidelberg, pp. 5–16.
- 4. Cooke, M.C. (1892) Vegetable Wasps and Plant Worms: A Popular History of Entomogenous Fungi, or Fungi Parasitic upon Insects. Society for Promoting Christian Knowledge, London.
- 5. Vega,F.E., Meyling,N.V., Luangsa-ard,J.J. et al. (2012) Fungal entomopathogens. Insect Pathol., 2, 171–220.
- Mora,M.A.E., Castilho,A.M.C. and Fraga,M.E. (2017) Classification and infection mechanism of entomopathogenic fungi. *Arq. Inst. Biol. (Sao Paulo)*, 84, 1–10.
- Gams, W. and Zare, R.A. (2003) Taxonomic review of the clavicipitaceous anamorphs parasitizing nematodes and other microinvertebrates. In: *Clavicipitalean Fungi*. pp. 26–81.
- 8. Bench, M.E. and White, M.M. (2012) New species and first records of Trichomycetes from immature aquatic insects in Idaho. *Mycologia*, 104, 295–312.
- 9. Araújo, J.P.M., Evans, H.C., Kepler, R. *et al.* (2018) Zombie-ant fungi across continents: 15 new species and new combinations within *Ophiocordyceps*. I. Myrmecophilous hirsutelloid species. *Stud. Mycol.*, **90**, 119–160.
- Hyde,K.D., Tennakoon,D.S., Jeewon,R. *et al.* (2019) Fungal diversity notes 1036–1150: taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Divers.*, 96, 1–242.
- Wei,D.P., Wanasinghe,D.N., Hyde,K.D. et al. (2020) Ophiocordyceps tianshanensis sp. nov. on ants from Tianshan mountains, R. P. China. Phytotaxa, 464, 277–292.
- Sung,G.H., Hywel-Jones,N.L., Sung,J.M. *et al.* (2007) Phylogenetic classification of Cordyceps and the clavicipitaceous fungi. *Stud. Mycol.*, 57, 5–59.
- Luangsa-ard, J.J., Tasanathai, K., Mongkolsamrit, S. et al. (2007) Atlas of Invertebrate-pathogenic Fungi of Thailand, Vol 1. BIOTEC, National Science and Tecnology Development Agency, Thailand.
- Kepler,R.M., Luangsa-Ard,J.J., Hywel-Jones,N.L. *et al.* (2017) A phylogenetically-based nomenclature for Cordycipitaceae (Hypocreales). *IMA Fungus*, 8, 335–353.
- Xiao, Y.-P., Hongsanan, S., Hyde, K.D. *et al.* (2019) Two new entomopathogenic species of *Ophiocordyceps* in Thailand. *MycoKeys*, 47, 53–74.
- Hyde,K.D., Norphanphoun,C., Maharachchikumbura,S.S.N. et al. (2020) Refined families of Sordariomycetes. Mycosphere, 11, 305–1059.
- Hammond, W. (1997) Laboulbeniales on beetles: host utilization patterns and species richness of the parasites. *Biodivers. Conserv.*, 6, 701–719.
- Tanada,Y. and Kaya,H.K. (1993) *Insect Pathology*. Academic Press. Inc. Harcourt Brace Avanovich, San Diego, pp. 359–361.
- Misra, J.K. and Horn, B.W. (eds). (2001) *Trichomycetes and Other Fungal Groups*. Science Publishers Inc, Enfield, New Hampshire, p. 396.
- Hibbett, D.S., Binder, M., Bischoff, J.F. et al. (2007) A higherlevel phylogenetic classification of the Fungi. Mycol. Res., 111, 509–547.
- James, T.Y., Kauff, F., Schoch, C.L. *et al.* (2006) Reconstructing the early evolution of fungi using a six-gene phylogeny. *Nature*, 443, 818–822.
- 22. Kirk,P.M., Cannon,P.F., Minter,D.W. et al. (2008) Stalpers, J.A. Ainsworth & Bisby's Dictionary of the Fungi. 10th edn. CAB International, Wallingford.
- Humber, R.A. (2012) Entomophthoromycota: a new phylum and reclassification for entomophthoroid fungi. *Mycotaxon*, 120, 477–492.

- 24. Spatafora, J.W., Chang, Y., Benny, G.L. *et al.* (2016) A phylum-level phylogenetic classification of zygomycete fungi based on genome-scale data. *Mycologia*, **108**, 1028–1046.
- Tedersoo,L., Sánchez-Ramírez,S., Kõljalg,U. *et al.* (2018) Highlevel classification of the Fungi and a tool for evolutionary ecological analyses. *Fungal Divers.*, 90, 135–159.
- Wijayawardene, N.N., Pawłowska, J., Letcher, P.M. *et al.* (2018) Notes for genera: basal clades of Fungi (including Aphelidiomycota, Basidiobolomycota, Blastocladiomycota, Calcarisporiellomycota, Caulochytriomycota, Chytridiomycota, Entomophthoromycota, Glomeromycota, Kickxellomycota, Monoblepharomycota, Mortierellomy. *Fungal Divers.*, 92, 43–129.
- 27. Martin, W.W. (1981) *Couchia circumplexa*, a water mold parasitic in midge eggs. *Mycologia*, 73, 1143–1157.
- 28. Mendoza,L., Vilela,R. and Humber,R.A. (2018) Taxonomic and phylogenetic analysis of the *Oomycota mosquito* larvae pathogen *Crypticola clavulifera*. *Fungal Biol.*, **122**, 847–855.
- 29. Vilela,R., Humber,R.A., Taylor,J.W. *et al.* (2019) Phylogenetic and physiological traits of oomycetes originally identified as *Lagenidium giganteum* from fly and mosquito larvae. *Mycologia*, **111**, 408–422.
- Charnley,A.K. and Collins,S.A. (2007) Entomopathogenic fungi and their role in pest control. In: Kubicek CP, Druzhinina, I. (eds.). *Environmental and Microbial Relationships: The Mycota IV*. 2nd edn. Springer-Verlag, Berlin Heidelberg, pp. 159–187.
- Boni,S.B., Mwashimaha,R.A., Mlowe,N. *et al.* (2021) Efficacy of indigenous entomopathogenic fungi against the black aphid, Aphis fabae Scopoli under controlled conditions in Tanzania. *Int. J. Trop. Insect Sci.*, 41, 1643–1651.
- 32. Mudrončeková,S., Mazáň,M., Nemčovič,M. et al. (2021) Entomopathogenic fungus species Beauveria bassiana (Bals.) and Metarhizium anisopliae (Metsch.) used as mycoinsecticide effective in biological control of Ips typographus (L.). J. Microbiol. Biotechnol. Food Sci., 2021, 2469–2472.
- Zhang,L., Fasoyin,O.E., Molnár,I. *et al.* (2020) Secondary metabolites from hypocrealean entomopathogenic fungi: novel bioactive compounds. *Nat. Prod. Rep.*, 37, 1181–1206.

- Maharachchikumbura, S.S., Chen, Y., Ariyawansa, H.A. et al. (2021) Integrative approaches for species delimitation in Ascomycota. Fungal Divers., 109, 155–179
- Raghoonundon,B., Zhao,R.-L., He,M.-Q. *et al.* (2021) www.basidio.org: an Online platform for the classification, identification and nomenclature of taxa within Basidiomycota. *Chiang Mai J. Sci.*, 48, 1186–1198.
- Wu,N., Dissanayake,A.J., Manawasinghe,I.S. *et al.* (2021) https://botryosphaeriales. org/, an online platform for up-to-date classification and account of taxa of Botryosphaeriales. *Database*, 2021, baab061.
- Pem,D., Hongsanan,S., Doilom,M. *et al.* (2019) https://www. dothideomycetes. org: an online taxonomic resource for the classification, identification, and nomenclature of Dothideomycetes. *Asian J. Mycol.*, 2, 287–297.
- Calabon,M.S., Hyde,K.D., Jones,E.B.G. *et al.* (2020) www. freshwaterfungi. org, an online platform for the taxonomic classification of freshwater fungi. *Asian J. Mycol.*, 3, 419–445.
- Jayawardena,R.S., McKenzie,E.H.C., Chen,Y.J. *et al.* (2019) https://onestopshopfungi. org/, a database to enhance identification of phytopathogenic genera. *Asian J. Mycol.*, 2, 281–286.
- Bundhun,D., Maharachchikumbura,S.S.N., Jeewon,R. *et al.* (2020) https://sordariomycetes. org/, a platform for the identification, ranking. *Asian J. Mycol.*, 3, 13–21.
- 41. Kobayasi,Y. (1941) The genus *Cordyceps* and its allies. *Rep. Tokyo Bunrika Daigaku Sect. B*, **5**, 53–260.
- 42. Kobayasi,Y. (1982) Keys to the taxa of the genera *Cordyceps* and *Torrubiella*. *Trans*. *Mycol*. *Soc*. *Japan*, 23, 329–364.
- 43. Mains, E.B. (1958) North American entomogenous species of *Cordyceps. Mycologia*, 50, 169–222.
- Massee,G. (1895) A revision of the genus Cordyceps. Ann. Bot., 9, 1–44.
- Mongkolsamrit,S., Khonsanit,A., Thanakitpipattana,D. et al. (2020) Revisiting Metarhizium and the description of new species from Thailand. Stud. Mycol., 95, 171–251.